

Epidemiology of 2,4-D Herbicide

Epidemiology studies address relationships of exposures and diseases in populations. The science of epidemiology provides important information about patterns of diseases in groups of people, such as pesticide applicators, farmers and farm workers. The process of identifying causes of disease within pesticide exposed populations is complex, mainly because pesticides are just one of many environmental exposures that people may encounter. Other exposures may include: fertilizers, nitrates, fuels, engine exhausts, solvents, organic and inorganic dusts, ultraviolet radiation, plant pathogens and animal pathogens. Behavioral, dietary and genetic factors may affect their risk of disease as well.

It is difficult to estimate exposure to humans since unlike controlled studies of laboratory animals, epidemiology studies can only approximate *potential* exposures. A common approach is to determine how many applications are made per year and the number of years applied. This approach can identify frequent and long term applicators from those who rarely apply 2,4-D. However, actual internal exposure may vary with acres applied, method of application and use of protective equipment; i.e., gloves, when mixing/loading. Detection levels in farmers' urine immediately after a 2,4-D application have demonstrated that exposure varies by individual and can be reduced by carefully following label instructions (Alexander 2007; Arbuckle 2002).

Exposure studies detect little more than trace levels of 2,4-D in urine of the general population, even though 2,4-D is one of the most commonly used herbicides in both the home and garden market and the commercial market (CDC 2015; Health Canada 2013) and children of applicators (Alexander 2007). Evaluations of these and other exposure studies have concluded that actual exposures are well below the conservative assumptions made by policy makers for the registration process (Hays 2012; Burns and Swaen 2012; Aylward and Hays 2015).

2,4-D Epidemiology Studies

There are numerous published epidemiology studies pertinent to 2,4-D. As highlighted by some non-government organizations, some of these studies looked at the links between 2,4-D and non-Hodgkin lymphoma (NHL) and soft-tissue sarcoma, both forms of cancer. These reports fail to clarify that early investigations – with weak methodology reported an association, but later independent investigations with more robust methods – did not validate these findings (Garabrant and Philbert 2001).

For example, two of these early studies are the National Cancer Institute's (NCI) studies in Kansas (Hoar 1986) and Nebraska (Zahm 1990), both of which received considerable media attention. A later NCI study in Iowa and Minnesota (Cantor 1992) showed no association between 2,4-D and NHL. The studies were critically

70
years
of and
RESEARCH
DISCOVERY

weakened by differential exposure perceptions among study respondents. All three studies relied heavily on information provided by proxy respondents.

That is, in many cases, the person chosen to participate in the study was not available, so next-of-kin or neighbors were asked about the unavailable person's pesticide use over a previous period of more than forty years.

Subsequent studies have shown that information on pesticide use provided by proxies is often invalid. In the NCI Nebraska study, for example, when the information provided by the self-respondents (i.e., the farmers themselves) is analyzed separately from the information provided by the neighbors or next-of-kin, there is no association between 2,4-D and cancer (Olsen 1996). The NCI combined the data from all three studies and reported, *"This analysis of the pooled data found no association with having ever used 2,4-D."* The weight of evidence from these NCI studies is that 2,4-D is not a carcinogen.

The extensive toxicology of 2,4-D (i.e., animal feeding studies done under controlled laboratory conditions in accordance with EPA GLP standards) does not support the hypothesis that 2,4-D is a carcinogen (WHO 1996; USEPA 2014; EFSA 2015), and the most robust studies in epidemiology consistently agree with the toxicology and biomonitoring studies. The review of the extensive toxicology and epidemiology of 2,4-D completed in 2001 by Garabrant and Philbert of the University of Michigan School of Public Health concluded:

"Despite several thorough in vitro and in vivo animal studies, no experimental evidence exists supporting the theory that 2,4-D or any of its salts or esters damages DNA under physiologic conditions. Studies in rodents demonstrate a lack of oncogenic or carcinogenic effects following lifetime dietary administration of 2,4-D. Epidemiologic studies provide scant evidence that exposure to 2,4-D is associated with soft tissue sarcoma, non-Hodgkin's lymphoma, Hodgkin's disease or any other cancer. Overall, the available evidence from epidemiologic studies is not adequate to conclude that any form of cancer is causally associated with 2,4-D exposure."

Recent published reviews and governmental regulatory decisions support the views of Garabrant and Philbert (von Stackelberg 2013; Burns and Swaen 2012; USEPA 2012; Health Canada PMRA 2008; USEPA 2005, EFSA 2014).

In 2015, Goodman et. al., conducted a meta-analysis of the published epidemiological literature for 2,4-D and concluded, "The epidemiology evidence does not support an association between 2,4-D and NHL, gastric cancer, or prostate cancer risk." The International Agency for Research on Cancer (IARC) utilized the

70
years
of and
RESEARCH
DISCOVERY

meta-analysis during the June 2015 Working Group and reported, “epidemiological studies did not find strong or consistent increases in risks of NHL or other cancers in relation to 2,4-D exposure.”

An excellent reference guide for further information on epidemiology studies is the Purdue University report titled, [“Pesticides and Epidemiology: Unraveling Disease Patterns”](#) (Purdue 1998).

Toxicology and epidemiology studies often referenced by those opposed to pesticides do not sufficiently document exposures and are inadequate to make cause and effect conclusions. In contrast, the extensive modern regulatory toxicology profile available for 2,4-D provides a robust and reassuring characterization of the compound. In conclusion, *“EPA has determined that this use is safe for humans and the environment when used according to the label.”* (USEPA 2014)

On October 14, 2014, the U.S. EPA published a memorandum in response to Public Comments received regarding an expanded 2,4-D use pattern in corn and soybeans. The agency concluded that:

“EPA thoroughly examined all of the significant environmental and health risks of 2,4-D including inhalation and aggregate exposure. EPA’s inhalation assessment evaluated risks using the most sensitive toxic effect – effects in the respiratory tract. Exposure to both residential pesticide users and professional applicators who would be applying the new 2,4-D formulation to herbicide resistant corn and soybeans were well below levels of risk concern.”

The agency goes on to state:

“EPA’s aggregate exposure assessment included contributions from food, drinking water, and non-occupational exposure, and was done for both adults and children. High-end, unrefined screening level inputs were used which resulted in exposure estimates well below levels of risk concern. Adult aggregate exposure included contributions from food, drinking water, and incidental oral exposure from swimming in a 2,4-D treated water body; the aggregate MOE was 1,800. Children’s aggregate exposure included contributions from food, drinking water, and incidental oral exposure hand to mouth from turf; the aggregate MOE was 340.”

70
 years
 of and
RESEARCH
DISCOVERY

About the Task Force

The Industry Task Force II on 2,4-D Research Data is organized to provide funding for the on-going Good Laboratory Practice (GLP) research studies required to respond to the US EPA registration review and PMRA pesticide re-evaluation programs. The 2,4-D Task Force is comprised of those companies holding technical 2,4-D registrations: Dow AgroSciences (U.S.), Nufarm, Ltd. (Australia) and Agro-Gor Corp., a U.S. corporation jointly owned by Albaugh, LLC. (U.S.) and PBI-Gordon Corp. (U.S.).

References:

Alexander, Bruce H., Jack S. Mandel, Beth A. Baker, Carol J. Burns, Michael J. Bartels, John F. Acquavella, and Christophe Gustin. (March 2007) Biomonitoring of 2,4-Dichlorophenoxyacetic Acid Exposure and Dose in Farm Families. *Environmental Health Perspectives*. Volume 115, No. 3: 370-376.

Arbuckle, Tye E., Burnett, R., Cole, D., Keschke, K., Dosemeci, M., Bacej, C. Zhang, J. (2002) Predictors of herbicide exposure in farm applicators. *Int Arch Occup Environ Health* 75: 406-414

Aylward, L.L., and S.M. Hays. (2015) Interpreting biomonitoring data for 2,4-dichlorophenoxyacetic acid: Update to Biomonitoring Equivalents and population biomonitoring data. *Regulatory Toxicology and Pharmacology* 73: 765-769.


Burns, Carol J. and Gerard M. H. Swaen. (2012) Review of 2,4-dichlorophenoxyacetic acid (2,4-D) biomonitoring and epidemiology. *Critical Reviews in Toxicology*. Vol. 42, No. 9:768-786.

Cantor, KP. et al. 1992. Pesticides and Other Agricultural Risk Factors for non-Hodgkin's Lymphoma Among Men in Iowa and Minnesota. *Cancer Research*, 52, 2447-2455.

CDC 2015. CDC Department of Health and Human Services: National Report on Human Exposure to Environmental Chemicals. Updated Tables, February 2015, Available: <http://www.cdc.gov/exposurereport/>.

De Roos, A.J. Zahm, SH. Cantor, KP. Weisenburger, DD. Holmes, FF. Burmeister, LF. and Blair, A. 2003. Integrative assessment of multiple pesticides as risk factors for non-Hodgkin's lymphoma among men. *Occup Environ Med*. 60(9): 1-9.

European Food Safety Authority. Conclusion on the peer review of the pesticide risk assessment of the active substance 2,4-D. EFSA Journal 2014;12(9):3812 (Scientific output, published on 11 March 2015, replaces the earlier version published on 7 August 2014)



Goodman, J.E., Loftus, C.T., Zu, K. 2,4-Dichlorophenoxyacetic acid and non-Hodgkin's lymphoma, gastric cancer, and prostate cancer: meta-analyses of the published literature. *Annals of Epidemiology* 25, 626–636. 2015.

Hays, S.M., Aylward, L.L., Driver, J., and Ross, J., Kirman, C. (2012) 2,4-D Exposure and risk assessment: Comparison of external dose and biomonitoring based approaches. *Summit Toxicology. Regulatory Toxicology and Pharmacology* 64:481-489.

Health Canada PMRA 2006. Re-evaluation Note REV2006-11 – August 2006. Lawn and Turf Uses of (2,4-Dichlorophenoxy)acetic Acid [2,4-D] Interim Measures.

Health Canada PMRA 2008. Re-evaluation Decision (2,4-Dichlorophenoxy) acetic Acid [2,4-D]. RVD2008-11.16 May 2008.

Health Canada 2013. Second Report on Human Biomonitoring of Environmental Chemicals in Canada, Results of the Canadian Health Measures Survey Cycle 2 (2009–2011). HC Pub.: 130019. Cat.: H128-1/10-601-1E-PDF. ISBN: 978-1-100-22140-3. Pages 262 – 269.

Hoar, SK. et. al. 1986. Agricultural Herbicide Use and Risk of Lymphoma and Soft-tissue Sarcoma, *JAMA*. 256:1141-7 see also correction, *JAMA* 256:3351.

Garabrant DH. Philbert MA. 2001 Review of 2,4-Dichlorophenoxyacetic Acid (2,4-D) Epidemiology and Toxicology. *CRC Critical Reviews in Toxicology*, 32(4):233-257.

International Agency for Research on Cancer, World Health Organization. Press Release No. 236 June 23, 2015

Olsen, GW. Bodner, KM. 1996. The Effect of the Type of Respondent on Risk Estimates of Pesticide Exposure in a Non-Hodgkin's Lymphoma Case-Control Study, *J. of Agromedicine*, Vol 3(1).

Purdue 1998. Purdue University Pesticide Programs, Report No. PPP-43, West Lafayette, Indiana 47907, July 1998.


USEPA 2005. Reregistration Eligibility Decision for 2,4-D. Office of Pesticide Programs. EPA 738-R-05-002. June 2005.

USEPA 2007. 2,4-D, 2,4-DP, and 2,4-DB; Decision Not to Initiate Special Review. Federal Register: August 8, 2007 (Volume 72, Number 152. Page 44510-44511)

USEPA 2012. Petitions to Revoke Tolerances; Denials: Natural Resources Defense Council, 2,4-dichlorophenoxyacetic acid. Federal Register 2012-09106. April 18, 2012 (Vol. 77, No. 75. Page 23135-23158)

USEPA 2014. Response to Public Comments Received Regarding New Uses of Enlist Duo™ on Corn and Soybeans", October 14, 2014.

USEPA 2014. Registration of Enlist Duo™ Herbicide Technical Briefing, October 15, 2014.



von Stackelberg, Katherine, 2013. A Systematic Review of Carcinogenic Outcomes and Potential Mechanisms from Exposure to 2,4-D and MCPA in the Environment. *Journal of Toxicology*, Volume 2013, Article ID 371610, 53 pages.

WHO Pesticide Residues in Food – 1996. Part II - Toxicological. FAO Plant Production and Protection Paper 140: 31-38; and WHO/PCS/97.1: 45-96.

Zahm, SH. et.al. 1990. Case-Control Study of Non-Hodgkin's Lymphoma and the Herbicide 2,4-Dichlorophenoxyacetic Acid (2,4-D) in Eastern Nebraska. *Epidemiology*, Vol 1, Number 5: 349-356.

March 17, 2016

70

years

of and

RESEARCH
DISCOVERY

INDUSTRY TASK FORCE II ON

2,4-D

RESEARCH DATA

24d.org